

1

"Cochlear implants in adults and children." NIH Consens Statement (1995) 13(2): 1-30.

2

Clark, G. M. "The development of the Nucleus multiple-channel cochlear implant." Aust J Otolaryng (1993) 1(4): 310-317.

3

Dawson, P. W. , Blamey, P. J. , Dettman, S. J. , Barker, E. J. ,Clark, G. M. "A clinical report on receptive vocabulary skills in cochlear implant users." Ear Hear (1995) 16: 287-294.

4

Dowell, R. C. , Blamey, P. J. ,Clark, G. M. "Potential and limitations of cochlear implants in children." Ann Otol Rhinol Laryngol Suppl (1995) 166: 324-327.

5

Eilers, R. E. , Cobo Lewis, A. B. , Vergara, K. C. , Oller, D. K. ,Friedman, K. E. "A longitudinal evaluation of the speech perception capabilities of children using multichannel tactile vocoders." J Speech Hear Res (1996) 39(3): 518-533.

6

Fryauf-Bertschy, H. , Tyler, R. S. , Kelsay, D. M. ,Gantz, B. J. "Performance over time of congenitally deaf and postlingually deafened children using a multichannel cochlear implant." J Speech Hear Res (1992) 35: 913-920.
(archive no: 683) -- in file Department of Otolaryngology-Head and Neck Surgery, University of Iowa, Iowa City 52242 The speech perception performance of 10 congenitally deaf and 3 postlingually deafened children who received the Cochlear Corporation multichannel cochlear implant was examined and compared. The children were tested preimplant and at 6-month intervals up to 2 years using the Monosyllable-Trochee-Spondee test (MTS), the Word Intelligibility by Picture Identification test (WIPI), and Phonetically Balanced Kindergarten (PB-K) or Northwestern University List 6 (NU-6) word lists. The postlingually deafened children exhibited significantly improved performance on open- and closed-set tests of word recognition after 6 months of implant use, a pattern similar to that of postlingually deafened adult implant users. In contrast, the congenitally deaf children did not exhibit measurably improved performance on speech perception tests until after 12 months or more of implant use. With as much as 18-24 months of use, however, some congenitally deaf children demonstrated limited open-set word recognition RefMgr field[1]: Journal RefMgr field[8]:

7

Geier, L. L. , Gilden, J. , Luetje, C. M. ,Maddox, H. E. "Delayed perception of cochlear implant stimulation in children with postmeningitic ossified cochleae." Am J Otol (1993) 14: 556-561.

8

Hodges, A. V. , Balkany, T. J. , Gomez-Marin, O. , Butts, S. , Ash, S. D. , Bird, P. ,Lee, D. "Speech recognition after implantation of the ossified cochlea." Am J Otol (1999) 20(4): 453-456.

9

Hodges, A. V. , Dolan, A. M. , Balkany, T. J. , Schloffman, J. J. ,Butts, S. L. "Speech perception results in children with cochlear implants: contributing factors." Otolaryngol Head Neck Surg (1999) 121(1): 31-34.

10

Kraus, N. , McGee, T. , Carrell, T. D. ,Sharma, A. "Neurophysiologic bases of speech discrimination." Ear Hear (1995) 16(1): 19-37.

11

Meyer, T. A. , Svirsky, M. A. , Kirk, K. I. ,Miyamoto, R. T. "Improvements in speech perception by children with profound prelingual hearing loss: effects of device, communication mode, and chronological age." J Speech Lang Hear Res (1998) 41(4): 846-858.

12

Miyamoto, R. T. , Kirk, K. I. , Svirsky, M. A. ,Sehgal, S. T. "Communication skills in pediatric cochlear implant recipients." Acta Otolaryngol (Stockh) (1999) 119(2): 219-224.

13

Miyamoto, R. T. , Osberger, M. J. , Robbins, A. M. , Myres, W. A. ,Kessler, K. S. "Prelingually deafened children's performance with the nucleus multichannel cochlear implant." Am J Otol (1993) 14: 437-445.
AD: Department of Otolaryngology-Head and Neck Surgery, Indiana University School of Medicine, Indianapolis 46202 AB: The speech perception abilities of 19 children with onset of deafness before age 3 years was examined after they received the Nucleus multichannel cochlear implant. The children were divided into two groups based on age at onset of deafness: children with congenital deafness (n = 8) and children with onset of deafness after birth but before age 3 (n = 11). There was no statistically significant difference between the scores of the two groups of subjects on 12 of the 13 speech perception tests administered. This finding suggests that children who are born deaf have the potential to derive the same benefit from cochlear implants as do children who have had some exposure to spoken language before the onset of their deafness. Examination of performance in terms of communication mode revealed that prelingually deafened children with implants who used oral communication obtained significantly higher scores on only 2 of the 13 speech perception measures than did children who used total communication. The data suggest that communication mode does not appear to account for large differences in speech perception performance among prelingually deafened children with multichannel cochlear implants RefMgr field[1]: Journal RefMgr field[8]:

14

Miyamoto, R. T. , Osberger, M. J. , Robbins, A. M. , Myres, W. A. , Kessler, K. S. ,Pope, M. L. "Longitudinal evaluation of communication skills of children with single- or multichannel cochlear implants." Am J Otol (1992) 13: 215-222.
Department of Otolaryngology-Head and Neck Surgery, Indiana University School of Medicine, Indianapolis Performance on speech perception, speech production, and language tests was measured over time in children who used the 3M/House or the Nucleus cochlear implant. The speech perception and production results demonstrated higher performance levels and faster rates of learning

for the multichannel than for the single-channel users. The performance of the children with the single-channel implant on the speech perception and production measures reached a plateau by 1.5 years post implant, whereas the children using the multichannel device continued to show improvement after 2 or more years of implant use. Changes in language were limited over time with no obvious difference in performance as a function of type of implant RefMgr field[1]: Journal RefMgr field[8]:

15

Miyamoto, R. T. , Osberger, M. J. , Todd, S. L. , Robbins, A. M. , Stroer, B. S. , Zimmerman Phillips, S. , Carney, A. E. "Variables affecting implant performance in children." *Laryngoscope* (1994) 104: 1120-1124.

16

Osberger, M. J. "Audiological rehabilitation with cochlear implants and tactile aids." *ASHA* (1990) 32: 38-43.

17

Osberger, M. J. , Miyamoto, R. T. , Zimmerman Phillips, S. , Kemink, J. L. , Stroer, B. S. , Firszt, J. B. , Novak, M. A. "Independent evaluation of the speech perception abilities of children with the Nucleus 22-channel cochlear implant system." *Ear Hear* (1991) 12: 66S-80S.

18

Osberger, M. J. , Zimmerman-Phillips, S. , Barker, M. , Geier, L. "Clinical trial of the CLARION cochlear implant in children." *Ann Otol Rhinol Laryngol Suppl* (1999) 177: 88-92.

19

Purdy, S. C. , Chard, L. L. , Moran, C. A. , Hodgson, S. A. "Outcomes of cochlear implants for New Zealand children and their families." *Ann Otol Rhinol Laryngol Suppl* (1995) 166: 102-105.

20

Robbins, A. M. , Svirsky, M. , Kirk, K. I. "Children with implants can speak, but can they communicate?" *Otolaryngol Head Neck Surg* (1997) 117(3 Pt 1): 155-160.

21

Roberts, S. A. , Busby, P. A. , Dowell, R. C. , Clark, G. M. , Tong, Y. C. , Mecklenburg, D. J. "Initial experience with children using the Nucleus 22 Channel cochlear implant." *Scand Audiol Suppl* (1988) 30: 115-119.

22

Snik, A. , Vermeulen, A. M. , Geelen, C. P. , Brokx, J. P. , van den Broek, P. "Speech Perception Performance of Children with a Cochlear Implant Compared to that of Children with Conventional Hearing Aids/Part II." *Acta Otolaryngol (Stockh)* (1997) 117(September): 755-759.

23

Somers, M. N. "Speech perception abilities in children with cochlear implants or hearing aids." *Am J Otol* (1991) 12 Suppl: 174-178.

24

Spencer, L. J. , Tye Murray, N. , Tomblin, J. B. "The production of English inflectional morphology, speech production and listening performance in children with cochlear implants." *Ear Hear* (1998) 19(4): 310-318.

25

Svirsky, M. A. , Meyer, T. A. "Comparison of speech perception in pediatric CLARION cochlear implant and hearing aid users." *Ann Otol Rhinol Laryngol Suppl* (1999) 177: 104-109.

26

Tye Murray, N. "Young cochlear implant users' response to delayed auditory feedback." *J Acoust Soc Am* (1992) 91: 3483-3486. Department of Otolaryngology--Head and Neck Surgery, University of Iowa Hospitals and Clinics, Iowa City 52242 This investigation determined whether the signal provided by the Cochlear Corporation Nucleus cochlear implant can convey enough speech information to induce a response to delayed auditory feedback (DAF), and whether prelingually deafened children who received a cochlear implant relatively late in their speech development are susceptible. Ten children with the Nucleus cochlear implant spoke simple phrases, first without and then with DAF. Three prelingually deafened subjects and the only two postlingually deafened subjects demonstrated longer phrase durations when speaking with DAF than without it. Two of the prelingually deafened subjects who demonstrated a response received their cochlear implants at the age of 5 years RefMgr field[1]: Journal RefMgr field[8]:

27

van den Borne, S. , Snik, A. F. , Hoekstra, C. C. , Vermeulen, A. M. , van den Broek, P. , Brokx, J. P. "Assessment of basal sound identification skills and communication abilities in profoundly deaf children fitted with hearing aids or a cochlear implant." *Clin Otolaryngol* (1998) 23(5): 455-461.

28

Vergara, K. C. , Oller, D. K. , Eilers, R. , Balkany, T. J. "Curricula objectives for educators of children with cochlear implants." *Adv Otorhinolaryngol* (1993) 48: 216-221. (archive no: 1259) -- in file AD: University of Miami, School of Medicine, Department of Pediatrics, Fla RefMgr field[1]: Journal RefMgr field[8]:

29

Vieu, A. , Mondain, M. , Blanchard, K. , Sillon, M. , Reuillard Artieres, F. , Tobey, E. , Uziel, A. , Piron, J. P. "Influence of communication mode on speech intelligibility and syntactic structure of sentences in profoundly hearing impaired French children implanted between 5 and 9 years of age." *Int J Pediatr Otorhinolaryngol* (1998) 44(1): 15-22.

30

Vormes, E. , Fugain, C. , Frachet, B. , Meyer, B. , Chouard, C. H. "[Presentation of a French language protocol for the evaluation of communication improvement in the deaf following implantation]." *Ann Otolaryngol Chir Cervicofac* (1990) 107: 466-468.

31

Waltzman, S. B. ,Cohen, N. L. "Cochlear implantation in children younger than 2 years old." *Am J Otol* (1998) 19(2): 158-162.

32

Waltzman, S. B. , Cohen, N. L. , Gomolin, R. H. , Green, J. E. , Shapiro, W. H. , Hoffman, R. A. ,Roland, J. T., Jr. "Open-set speech perception in congenitally deaf children using cochlear implants." *Am J Otol* (1997) 18: 342-349.

33

Zeng, F. G. "Cochlear implants in China." *Audiology* (1995) 34(2): 61-75.